

09/509, 775



CGCTCAGGATACGACTTCGGCTAGAGATCGGGATCCCGGCGGATTATTATAGCTCGATCGATCT  
TTCTCTATATCCGGCGGATGGGATATATACACACACACCGCGCGATAGCATGACTGATCTA  
CCCCATCT  
CAGAGACTCTACGCT

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

B

Search

Nucleotide



for gankyrin

Preview

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Search for Genes  
LocusLink provides  
curated information for  
human, fruit fly, mouse,  
rat, and zebrafish

- Search History will be lost after one hour of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Search

Most Recent Queries

Time Result

Entrez Nucleotide

#1 Search gankyrin

14:52:53

40

Help | FAQ

Batch Entrez: Upload a  
file of GI or accession  
numbers to retrieve  
sequences

Clear History

Check sequence  
revision history

How to create WWW  
links to Entrez

LinkOut

Cubby

Related resources  
BLAST

Reference sequence  
project


Submit to GenBank

Revised: July 5, 2002.

Disclaimer | Write to the Help Desk  
NCBI | NLM | NIH

i686-pc-linux-gnu Jan 21 2003 17:57:06

**OMIM**  
*Online Mendelian Inheritance in Man*

 **Johns  
Hopkins  
University**

PubMed   Nucleotide   Protein   Genome   Structure   PMC   Taxonomy   OMIM

Search  for


Limits   Preview/Index   History   Clipboard   Details

Show:

**\*603480**

Links

**PROTEASOME 26S SUBUNIT, NON-ATPase, 10; PSMD10****Alternative titles; symbols****p28****TEXT**

Ubiquitinated proteins are degraded by a 26S ATP-dependent protease. The protease is composed of a 20S catalytic proteasome and PA700, a 700-kD regulatory complex (see PSMC1, 602706). Hori et al. (1998) determined the partial protein sequences of bovine p28 and p40.5 (PSMD13; 603481), 2 components of PA700. By searching a sequence database, they identified cDNAs encoding the human p28 and p40.5 homologs. The predicted 226-amino acid human p28 protein contains 5 ankyrin repeats, which are thought to function in protein-protein interactions. Using computerized homology searches, Hori et al. (1998) identified Nas6, an *S. cerevisiae* gene encoding a protein with 38% identity to p28. They found that disruption of Nas6 had no effect on cell viability. Northern blot analysis revealed that p28 was expressed as a 1.3-kb mRNA in all human tissues tested. 

**REFERENCES**

1. Hori, T.; Kato, S.; Saeki, M.; DeMartino, G. N.; Slaughter, C. A.; Takeuchi, J.; Toh-e, A.; Tanaka, K. :  
cDNA cloning and functional analysis of p28 (Nas6p) and p40.5 (Nas7p), two novel regulatory subunits of the 26S proteasome. *Gene* 216: 113-122, 1998.  
PubMed ID : [9714768](#)

**CREATION DATE**

Rebekah S. Rasooly : 2/3/1999


**EDIT HISTORY**

alopez : 2/3/1999

Copyright © 1966-2003 Johns Hopkins University

|         |          |       |    |         |      |
|---------|----------|-------|----|---------|------|
| Display | Detailed | Show: | 20 | Send to | Text |
|---------|----------|-------|----|---------|------|

[Disclaimer](#) | [Write to the Help Desk](#) | [Privacy Policy](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[PMC](#)[Taxonomy](#)[OMIM](#)[B](#)

Search

PubMed

for

Go

Clear

[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)[About Entrez](#)

Display

Abstract

Show:

20

Sort

Send to

Text

[Text Version](#)**Entrez PubMed**[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)**PubMed Services**[Journals Database](#)[MeSH Browser](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[LinkOut](#)[Cubby](#)**Related Resources**[Order Documents](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)[Privacy Policy](#)☐ 1: Gene 1998 Aug 17;216(1):113-22[Related Articles, Links](#)**ELSEVIER SCIENCE  
FULL-TEXT ARTICLE**

**cDNA cloning and functional analysis of p28 (Nas6p) and p40.5 (Nas7p), two novel regulatory subunits of the 26S proteasome.**

Hori T, Kato S, Saeki M, DeMartino GN, Slaughter CA, Takeuchi J, Toh-e A, Tanaka K.

The Tokyo Metropolitan Institute of Medical Science, and CREST, Japan Science and Technology Corporation (JST), 3-18-22 Honkomagome, Bunkyo-ku, Tokyo 113-8613, Japan.

We employed cDNA cloning to deduce the complete primary structures of p28 and p40.5, two novel subunits of PA700 (also called 19S complex), a 700 kDa multisubunit regulatory complex of the human 26S proteasome. These polypeptides consisted of 226 and 376 amino acids with calculated molecular masses of 24428 Da and 42945 Da, and isoelectric points of 5.68 and 5.46, respectively. Intriguingly, p28 contained five conserved motifs known as 'ankyrin repeats', implying that this subunit may contribute to interaction of the 26S proteasome with other protein(s). Computer-assisted homology analysis revealed high sequence similarities of p28 and p40.5 with yeast proteins, termed Nas6p and Nas7p (non-ATPase subunits 6 and 7), respectively, whose functions are as yet unknown. Disruption of these yeast genes, NAS6 and NAS7, had no effect on cell viability, indicating that neither of the two subunits is essential for proliferation of yeast cells. However, the NAS7, but not NAS6, disruptant cells caused high sensitivity to heat stress, being unable to proliferate at 37 degreesC.

PMID: 9714768 [PubMed - indexed for MEDLINE]